

Necessity of Dark Matter in Modified Newtonian Dynamics within Galactic Scales? - Testing the Covariant MOND in Elliptical Lenses

M.C. Chiu¹, Y. Tian² and C.M. Ko³

¹*Institute of Astronomy, National Central University, Taiwan*

²*Department of Physics, National Central University, Taiwan*

³*Institute of Astronomy, Department of Physics and Center of Complex System, National Central University, Taiwan*

ABSTRACT

Modified Newtonian Dynamics (MOND) and its relativistic version – TeVeS offer us an alternative perspective to understand the universe without the demand of the elusive cold dark matter. This MONDian paradigm is not only competitive with the conventional CDM in a large range of scales, but also even more successful in the galactic scale. Recently, by studying 6 lensing systems, Ferreras et al. (2008) claimed that MOND still needs dark matter even in galactic scales. When we study the same systems, however, we yield an opposite conclusion. In this contribution, we report our result and conclude that MOND does not need dark matter in galactic lensing systems. Furthermore, we extend our study to 22 SLACS (Sloan Lens ACS Survey) lenses, and obtain the same conclusion as well, i.e., no dark matter is needed in elliptical galaxies.

Key words: Gravitational lensing - MOND - dark matter - gravitation - relativity

1 INTRODUCTION

Migrom’s M^ODified Newtonian Dynamics (MOND) is an alternative to the conventional dark matter paradigm for the rotation curve of galaxies and similar phenomena (Migrom 1983). TeVeS-a relativistic MOND theory (Bekenstein 2004) offered an opportunity to explain relativistic phenomena such as cosmology and gravitational lensing (Chiu et al. 2006).

Recently, Ferreras et al. (2008) studied 6 lensing systems and claimed that MOND still needs dark matter even in galactic scales. We revisit the problem and arrive at an opposite conclusion. Here we report our result on 10 lens from CASTLES (including the 6 lens studied by Ferreras et al.) and 22 lens from SLACS.

2 LENSING EQUATION

Since in the MONDian paradigm mass distribution only follows baryon, we adopt the Hernquist model, $|\nabla\Phi_N| = GM/(r+r_h)^2$, to the lenses. We incorporate our lensing formalism in a ν HDM cosmological background ($\Omega_b = 0.05$, $\Omega_\nu = 0.17$ $\Omega_\Lambda = 0.78$, $h = 0.7$).

In general, for any double-images lensing systems:

$$\theta_+\theta_- = \theta_E^2 \left(\frac{\theta_+f_- + \theta_-f_+}{\theta_+ + \theta_-} \right), \quad (1)$$

where $\theta_\pm$ are positions of the two images, $\theta_E^2 = 4GMD_{LS}/c^2D_LD_S$, and $f_\pm$ are dimensionless functions, which depend on the mass model of lenses and the forms of $\tilde{\mu}(x)$ in MOND. We consider (i) the Bekenstein’s form $\tilde{\mu} = (-1 + \sqrt{1+4x})/(1 + \sqrt{1+4x})$ (Bekenstein 2004), (ii) the simple form $\tilde{\mu} = x/(1+x)$ (Famaey & Binney 2005), and (iii) the standard form $\tilde{\mu} = x/\sqrt{1+x^2}$ (Sanders & McGaugh 2002). Here $x = a/a_0$, the ratio of the actual acceleration to the acceleration parameter in MOND.

3 DATA AND RESULT

We apply 10 CASTLES lensing systems along with 22 SLACS lenses to study strong lensing in TeVeS.


CASTLES Catalogue has the most complete list of strong lensing systems. We examine 10 double-image lenses from the this catalogue, of which the (aperture) stellar masses have been estimated from stellar population synthesis with two initial mass functions (IMF), Salpeter’s and Chabrier’s (Ferreras et al. 2005). In Table 2 we list the aperture mass and total mass computed from the three forms of $\tilde{\mu}(x)$ (Bekenstein, simple and standard).

We also work out the masses for 22 SLACS lens. SLACS lens come from SDSS Luminous Red Galaxy and MAIN SDSS galaxy sample, in which the dispersion velocity can be estimated. We use the M- σ relation in (Sanders 2000) to create a M- σ -R_{eff} relation of V-band in MOND, and apply

Table 1. Aperture mass (total mass) of 10 lenses from CASTLES ($10^{10} M_{\odot}$) in ν HDM

Lens	Bekenstein	Simple	Standard
Q0142 – 100	10.79 (18.36)	13.66 (23.20)	16.05 (27.31)
HS0818 + 1227	18.14 (28.30)	23.39 (36.17)	27.79 (43.35)
FBQ0951 + 2635	1.54 (2.16)	1.91 (2.67)	2.15 (3.01)
BRI0952 – 0115	2.01 (2.48)	2.59 (3.21)	3.19 (3.93)
Q1017 – 207	2.45 (5.89)	3.22 (7.55)	3.81 (9.15)
HE1104 – 1805	45.17 (59.58)	58.44 (77.08)	71.78 (94.68)
LBQ1009 – 025	7.71 (10.79)	9.76 (13.67)	11.53 (16.15)
B1030 + 071	9.76 (16.61)	12.06 (20.51)	13.80 (23.47)
SBS1520 + 530	11.91 (16.67)	15.20 (21.28)	18.08 (25.31)
HE2149 – 274	7.04 (13.58)	8.96 (17.28)	10.67 (20.58)

Lens	Chabrier	Salpeter
Q0142 – 100	20.9 ^{30.8} _{13.0}	18.3 ^{32.2} _{13.2}
HS0818 + 1227	16.2 ^{21.2} _{12.6}	20.8 ^{28.1} _{13.4}
FBQ0951 + 2635	1.1 ^{1.1} _{0.5}	1.5 ^{3.0} _{0.8}
BRI0952 – 0115	3.5 ^{4.0} _{1.7}	4.4 ^{5.0} _{3.5}
Q1017 – 207	4.3 ^{13.0} _{1.4}	6.4 ^{19.0} _{2.3}
HE1104 – 1805	22.8 ^{51.2} _{12.7}	36.6 ^{63.7} _{23.1}
LBQ1009 – 025	5.5 ^{7.9} _{4.2}	7.4 ^{9.8} _{5.0}
B1030 + 071	10.6 ^{15.3} _{6.5}	14.5 ^{21.3} _{8.3}
SBS1520 + 530	18.5 ^{30.9} _{11.2}	21.8 ^{34.1} _{11.9}
HE2149 – 274	4.6 ^{6.7} _{3.6}	6.9 ^{9.9} _{5.0}

**Figure 1.** Comparison of mass from lensing and mass from dynamical measurement (i.e., velocity dispersion). Simple form gives the best correlation.

this formula to compute the dynamical mass. The comparison between lensing mass and dynamical mass is given in Table 2. (θ/θ_0) is a measure of how close the lensing system is to the deep MOND or the Newtonian regime. All the 22 systems are in the intermediate MOND regime, in which the form of $\tilde{\mu}(x)$ is important. In Fig. 1, we compare the mass from lensing and mass from velocity dispersion.

4 SUMMARY

Our investigation supports that there is no need of dark matter in elliptical galaxies.

Among the three choices of $\tilde{\mu}(x)$ of MOND, simple form seems to yield the best fit with dynamical counterparts. This echoes the dynamical studies in spiral galaxies (Famaey & Binney 2005; Famaey et al. 2007).

ACKNOWLEDGMENT

This work is supported in part by NSC96-2112-M-008-014-MY3.

Table 2. Mass of 22 lenses from SLACS ($10^{10} M_{\odot}$) in ν HDM

Lens	$(\frac{\theta}{\theta_0})_{\text{bek}}$	$(\frac{\theta}{\theta_0})_{\text{spl}}$	$(\frac{\theta}{\theta_0})_{\text{std}}$
SDSS J002907.8 – 005550	0.227	0.200	0.183
SDSS J015758.9 – 005626	0.338	0.298	0.274
SDSS J021652.5 – 081345	0.152	0.135	0.126
SDSS J025245.2 + 003958	0.246	0.216	0.197
SDSS J033012.1 – 002052	0.304	0.267	0.243
SDSS J072805.0 + 383526	0.247	0.218	0.203
SDSS J080858.8 + 470639	0.253	0.224	0.205
SDSS J090315.2 + 411609	0.301	0.263	0.240
SDSS J091205.3 + 002901	0.113	0.101	0.095
SDSS J095944.1 + 041017	0.164	0.146	0.137
SDSS J102332.3 + 423002	0.242	0.214	0.199
SDSS J110308.2 + 532228	0.138	0.121	0.112
SDSS J120540.4 + 491029	0.181	0.161	0.150
SDSS J125028.3 + 052349	0.241	0.212	0.195
SDSS J140228.1 + 632133	0.189	0.167	0.156
SDSS J142015.9 + 601915	0.111	0.102	0.097
SDSS J162746.5 – 005358	0.181	0.160	0.149
SDSS J163028.2 + 452036	0.264	0.232	0.213
SDSS J223840.2 – 075456	0.178	0.158	0.148
SDSS J230053.2 + 002238	0.180	0.160	0.149
SDSS J230321.7 + 142218	0.163	0.145	0.135
SDSS J234111.6 + 000019	0.183	0.161	0.148

Lens	M_{bek}	M_{spl}	M_{std}	M_{σ}
SDSS J002907.8 – 005550	15.22	19.68	23.45	36.8 ^{42.0} _{31.8}
SDSS J015758.9 – 005626	29.68	38.12	45.13	77.9 ^{101.1} _{57.4}
SDSS J021652.5 – 081345	119.91	151.50	172.79	126.9 ^{142.8} _{111.9}
SDSS J025245.2 + 003958	25.84	33.44	40.27	24.9 ^{28.2} _{21.8}
SDSS J033012.1 – 002052	27.26	35.43	42.76	35.0 ^{41.3} _{29.1}
SDSS J072805.0 + 383526	25.72	32.95	38.35	29.4 ^{32.0} _{26.7}
SDSS J080858.8 + 470639	26.01	33.38	39.57	40.9 ^{44.3} _{37.6}
SDSS J090315.2 + 411609	41.75	54.77	65.92	44.2 ^{54.1} _{35.2}
SDSS J091205.3 + 002901	139.99	176.47	197.36	126.7 ^{135.0} _{118.6}
SDSS J095944.1 + 041017	16.17	20.40	23.24	24.4 ^{27.3} _{21.6}
SDSS J102332.3 + 423002	25.48	32.43	37.66	35.9 ^{39.9} _{32.1}
SDSS J110308.2 + 532228	23.91	30.93	36.23	36.8 ^{40.8} _{32.9}
SDSS J120540.4 + 491029	35.24	44.65	51.29	59.9 ^{64.8} _{55.1}
SDSS J125028.3 + 052349	27.77	35.99	42.20	47.3 ^{52.0} _{42.7}
SDSS J140228.1 + 632133	53.51	68.51	78.22	59.6 ^{66.5} _{54.2}
SDSS J142015.9 + 601915	10.24	12.21	13.35	19.3 ^{21.7} _{12.7}
SDSS J162746.5 – 005358	45.52	58.40	67.44	72.9 ^{79.2} _{66.8}
SDSS J163028.2 + 452036	63.71	82.47	97.16	64.9 ^{71.6} _{58.4}
SDSS J223840.2 – 075456	22.68	28.72	32.96	28.3 ^{31.1} _{25.6}
SDSS J230053.2 + 002238	56.99	72.55	83.09	67.1 ^{74.5} _{60.1}
SDSS J230321.7 + 142218	62.56	79.12	90.74	62.9 ^{70.0} _{56.1}
SDSS J234111.6 + 000019	41.77	53.97	63.75	44.2 ^{49.2} _{39.5}

REFERENCES

- Bekenstein J.D., 2004, PRD 70, 083509
 Chiu M.C., Ko C.M., Tian Y. ApJ, 636, 565
 Famaey B., Binney J., 2005, MNRAS 363, 603
 Famaey B., Gentile G., Bruneton J.-P., Zhao H.S., 2007, PRD 75, 063002
 Ferreras I., Saha P., Williams L., 2005, ApJ 623, L5
 Ferreras I., Sakellariadou M., Yusaf M.F., 2008, PRL 100, 031302
 Milgrom M., 1983, ApJ 270, 365
 Sanders R.H., 2000, MNRAS 313, 767
 Sanders R.H., McGaugh S.S., 2002, ARAA 40, 263